## "WHAT DOES A WOUND FROM TRANQDOPE LOOK LIKE?"

TRING DOVE

WHAT YOU SHOULD KNOW ABOUT CARING FOR YOUR WOUNDS

BOUGHT TO YOU BY
FLORIDA HARM REDUCTION
COLLECTIVE'S COMMUNITY
ADVISORY BOARD. CHECK
US OUT FLHRC.ORG

Wounds can look like a burn or an ulcer. At its beginning, it may be red and swollen, yellow and goopy. They may also be black (dead tissue), dark purple, and flakey. Because xylazine constricts veins and slows down blood flow, you can get wounds that aren't at the IV site. Healthy bloodflow is not getting to previous wounds. This means, even if you are only smoking or snorting dope, you can end up with this type of wound.


STEP 1: Wash wound with soap and water (peroxide and alcohol are too harsh)

**STEP 2**: Keep wound moist- this helps dead skin to soften and fall off on its own. Xeroform bandages, or vaseline work well. Remember to cover it w/ non-adhesive absorbent pad/gauze after moisturizing it, then wrap with ace bandage.

STEP 3: Repeat process every 1-2 days.

PLEASE NOTE: We do not know a whole lot about the effects of tranqdope. This info is not a substitute for medical advice. If you have symptoms of a fever, your wound smells bad, or is extremely painful and getting worse- it's best to seek medical care!





1. ADMINISTER NARCAN & WAIT

Once they are breathing again, they may still be unconscious. Put them in the recovery position. More Narcan won't wake them up. If you're able to, call HMS, if not, stay with them for 2-3 hrs and count their breaths- 10-12 breaths per minute. If it drops below 10 breaths per minute, administer more Narcan and give rescue breaths. If they are consistently breathing 10 breaths per min, stay with them until they wake up.

PUT THEM IN RECOVERY

POSITION IF YOU HAVE TO